



## Research Article

### PHARMACOGNOSY OF PEETHA BHRINGARAJA (*WEDELIA CHINENSIS* (OSBECK) MERRILL)

Krishna Priya. A<sup>1\*</sup>, Shincymol. V.V<sup>2</sup>

\*1PG scholar, <sup>2</sup>Associate Professor, Department of Dravyaguna vijnanam, Govt. Ayurveda College, Tripunithura, Kerala, India.

#### ABSTRACT

*Bhringaraja* (*Eclipta prostrata* (L.)) is a plant widely used as a remedy for liver disorders, Anaemia etc. The drug is said to have three varieties based on the colour of its flower. *Peetha Bhringaraja* is the yellow flowered variety of *Bhringaraja*. The drug is botanically identified as *Wedelia chinensis* (Osbeck) Merrill belongs to the family Asteraceae. The drug is not that much abundantly seen nowadays. *Wedelia trilobata* is another similar invasive species belongs to the same family Asteraceae. The IUCN has listed *Wedelia Trilobata* in its 100 world's worst invasive alien species. Florida exotic plant pest council considered it as category 2 invader. Due to its invasion most of the similar *Wedelia* species got replaced. Pharmacognosy is the only reliable tool to differentiate among plants. For the purpose of utilisation of genuine source of *Wedelia chinensis* (Osbeck) Merrill, the plant was identified and detailed macroscopy and microscopy of root, stem and leaf along with the powder microscopy of whole plant is done.

**KEYWORDS:** *Peetha Bhringaraja*, *Wedelia chinensis* (Osbeck) Merrill, Macroscopy, Microscopy, Powder microscopy.

#### INTRODUCTION

*Peetha Bhringaraja* is the yellow coloured variety of *Bhringaraja*. It is *Vathakapha Hara* and does *karmas* like *Muthrala*, *Hridya*, *Vrishya*, *Swedakara*, *Kesya*, *Balya*.<sup>[1]</sup> It is botanically identified as *Wedelia chinensis* (Osbeck) Merrill belongs to the family Asteraceae. It is a procumbent perennial herb with stem rooting at nodes, growing upto 30cm to 1m in height.<sup>[2]</sup> The drug is often mistaken with *Wedelia trilobata*, which is another invasive species of same family. Pharmacognosy is the tool for identifying genuine drug. Pharmacognostical evaluation of a drug includes macroscopical (organoleptic) evaluation, microscopical evaluation and powder microscopy.

#### MATERIALS AND METHOD

**Collection of sample:** The whole plant of *Wedelia chinensis* (Osbeck) Merrill is collected from its natural

#### RESULT

Macroscopical or organoleptic evaluation of each part of the plant is done and it is tabulated.

habitat, washed thoroughly to remove soil and other impurities.

#### Method of study

Root, stem and leaf of the plant is studied macroscopically by evaluating its organoleptic features. For microscopical evaluation, thin section of each part is taken using blade and stained with saffranine, placed over a clean glass slide which is covered with a cover slip and observed under microscope. After analysing the sections in different magnifications different cells are identified. For doing powder microscopy, the whole plant of the drug which is cleaned well was dried and powdered. A pinch of powder is kept in a clean dry glass slide and a drop of glycerin was added to it and a thin smear is prepared and dried. This is then observed under microscope and cells are identified.



Figure 1: Whole plant



Figure 2: Root


**Root of *Peetha Bhringaraja* [*Wedelia chinensis* (Osbeck) Merrill]**

**Table 1: Organoleptic Evaluation**

| Parameters        | Root of <i>Peetha Bhringaraja</i> [ <i>Wedelia chinensis</i> (Osbeck) Merrill] |
|-------------------|--|
| Shape             | Narrow, elongated, rooting at nodes with lateral rootlets                      |
| Size              | Varying  |
| Colour            | Pale creamish /greyish to Buff / greyish brown in colour                       |
| Texture           | Nodular at some areas, Possess lateral rootlets                                |
| Odour             | Not characteristic   |
| Taste             | Not characteristic   |
| Fracture          | Short  |
| Fractured surface | Light creamish colour  |



**Figure 3: T.S of root (10X) showing outer cork, cortex and large air space**



**Figure 4: T.S of root (10X) showing outer cork, cortex, stone cells in cortex and not well developed pith**

**Stem of *Peetha Bhringaraja* [*Wedelia chinensis* (Osbeck) Merrill]**




**Figure 5: Stem**


**Table 2: Organoleptic Evaluation of Stem**

| Parameters          | Stem of <i>Peetha Bhringaraja</i> [ <i>Wedelia chinensis</i> (Osbeck) Merrill] |
|---------------------|--|
| Shape               | Cylindrical, prominent or bulged at nodes                                      |
| Colour              | Reddish brown to brown in colour   |
| External characters | Presence of whitish appressed hairs  |
| Texture             | Rough due to presence of hairs   |
| Odour               | Not characteristic   |
| Taste               | Slightly bitter  |
| Fracture            | Slightly fibrous   |


**Microscopy of stem**



**Figure 6: T.S of Stem (4X) showing outer Epidermis, cortex with air spaces, vascular bundles and pith**



**Figure 7: T.S of Stem (10X) showing outer Epidermis, cortex, air space, vascular bundles and pith**



**Figure 8: Vascular bundles 40X**

**Leaf of Peetha Bhringaraja [*Wedelia chinensis* (Osbeck) Merrill]**



**Figure 9: Leaf**


**Table 3: Organoleptic evaluation of leaf**

| Parameters | Leaf of <i>Peetha Bhringaraja (Wedelia chinensis (Osbeck) Merrill)</i> |
|------------|--|
| Kind       | Simple   |
| Size       | 2.5 – 7.5cm long and 7mm – 3.2cm wide                                  |
| Petiole    | Short, Pale coloured   |
| Shape      | Linear – oblong or oblanceolate or oblong lanceolate                   |
| Base       | Tapering   |
| Lamina     | Triple nerved  |
| Apex       | Acute  |
| Margin     | Entire / irregularly sub crenate, or crenate – serrate                 |
| Surface    | Rough due to white appressed hairs                                     |
| Colour     | Green  |
| Texture    | Rough  |
| Odour      | Not characteristic   |
| Taste      | Characteristic   |
| Fracture   | Brittle  |

**Microscopy of leaf**




**Figure 10: T.S of Midrib portion of leaf (10X) showing cuticle, epidermis, spongy parenchyma and vascular tissue in the centre**



**Figure 11: T.S through lamina of leaf (40X) showing palisade parenchyma**



**Figure 12: Warty trichome (40X)**



**Figure 13: Stomata (40X)**



**Figure 14) Glandular trichome (40X)**

**Powder Analysis**

**Table 4: Organoleptic evaluation of Powder**

| Parameters          | Powder of whole plant of <i>Peetha Bhringaraja</i> |
|---------------------|--|
| Colour              | Greenish grey                                      |
| External appearance | Fibrous  |
| Odour               | Characteristic                                     |
| Taste               | Slight bitter and astringent                       |

**Powder Microscopy**



**Figure 15: Powder of whole plant of *Peetha Bhringaraja***



**Figure 16: Lignified fibre**




**Figure 17: Pitted vessel**



**Figure 18: Spiral vessel**




**Figure 19: Trichome**



**Figure 20: Parenchymal cells**



**Figure 21: Cut fragment of lamina of leaf**



**Figure 22: Cork**



**Figure 23: Starch grain**

## CONCLUSION

Pharmacognostical analysis of different parts of the drug *Wedelia chinensis* (Osbeck) Merrill was done. The root and stem of the drug was mainly characterised by large air cells in the cortex region. Leaf contains anisocytic type of stomata. Characteristic of all part was evident from the powder analysis. From the details pharmacognostical evaluation the drug *Wedelia chinensis* (Osbeck) Merrill can be differentiated from other *Wedelia* species.

## ACKNOWLEDGEMENT

I am greatly thankful to Dr.P.Y.Ansary, Professor and H.O.D, Department of Dravyaguna vijnanam, GAVC, Tripunithura for his constant guidance and motivation. I am thankful to

Dr.Shincymol.V.V, my guide, for her immense support and guidance throughout the work. I express my deep gratitude to Dr.Sara Moncy Oommen, for her moral support and advices.

## REFERENCES

1. Ayurveda Pharmacopoeia of India, 1<sup>st</sup> edition, Government of India, Ministry of Health and Family welfare, Part 1,Vol 6.
2. M.Kolammal,Prof.K.NarayanaIyer, Pharmacognosy of Ayurvedic drugs, Trivandrum, Dept. of Pharmacognosy University of Kerala, Series 1, Number 5, P. 122.
3. A.K.Gupta, Quality standards of Indian medicinal plants, Volume 1, ICMR, 2003, P. 227.

### Cite this article as:

Krishna Priya. A, Shincymol. V.V. Pharmacognosy of Peetha Bhringaraja (*Wedelia Chinensis* (Osbeck) Merrill). International Journal of Ayurveda and Pharma Research. 2019;7(7):17-22.

**Source of support: Nil, Conflict of interest: None Declared**

### \*Address for correspondence

**Dr. Krishna Priya. A**

PG Scholar,  
Department of Dravyagunavijnana,  
Govt. Ayurveda College, Dhanwanthari  
nagar, Puthiyakavu, Tripunithura- 682301  
Phone: 9567418372  
Email: [drkrishnapriya44@gmail.com](mailto:drkrishnapriya44@gmail.com)

Disclaimer: IJAPR is solely owned by Mahadev Publications - dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IJAPR cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of IJAPR editor or editorial board members.

